



RADT Vs RT-PCR

Why in News

Recently, New Delhi's [Covid-19](#) testing strategy has become controversial due to **the low level of RT-PCR** (Reverse Transcription Polymerase Chain Reaction) **re-testing in persons** tested negative in **RADT** (Rapid Antigen Detection Tests).

- Using RADT **widely without following up** with **adequate retests contradicts** [Indian Council of Medical Research \(ICMR\) guidelines](#) on use of the RADT test.

Key Points

▪ ICMR Guidelines:

- RADT ought to be used only in **containment zones, hotspots, hospital settings and laboratories** among those who manifested one or other symptoms of the disease, influenza-like illnesses.
- People with **comorbidities who were asymptomatic** and high-risk contacts of those confirmed positive.
- Those who tested 'negative' and whom **clinicians suspected** to be harbouring the disease ought to be **definitely tested sequentially by RT-PCR** to rule out infection and **higher chances of false negatives**.
- Those who test positive don't need a re-test and must be considered positive.

▪ Testing in New Delhi:

- From 18th June - 16th July, it has conducted 3,05,820 RADT. Of these, 2,85,225 tests came 'negative' and out of them only, 1,670 were chosen for re-test by RT-PCR and 262 of these were confirmed positive.
- Only **1 in 200 of those who tested negative in an antigen** test to detect possible coronavirus cases **were re-tested**, which is against the given guidelines of ICMR.
- Of those re-tested with **RT-PCR, around 15% tested positive**, which is higher than the **RADT positive results i.e. 6%**.

▪ Arguments for Low Re-tests:

- Re-testing everyone would defeat the purpose of having another (rapid antigen) test.
- The RT-PCR test takes a minimum of 2-5 hours including the time taken for sample transportation. This **limits the widespread use of the test** and also impedes quick augmentation of testing capacity in various containment zones and hospital settings.

- In **RADT, the maximum duration** for interpreting a positive or negative test is **30 minutes**, thus **a quicker complement to the standard RT-PCR** tests.

▪ Arguments Against:

- The consequence of **indiscriminately deploying antigen tests** would mean expanding the number of tests and presenting a lower positivity rate while not necessarily being able to **reliably establish the extent of the spread** of the coronavirus in the population.
- A low level of re-testing with RT-PCR in persons who are testing antigen negative will

underestimate the cases and make the **tracking inaccurate**.

RADT

- It is a test on swabbed nasal samples that detects antigens (foreign substances that induce an immune response in the body) that are found on or within the SARS-CoV-2 virus.
- It is a **point-of-care test**, performed outside the conventional laboratory setting, and is used to quickly obtain a diagnostic result.
- Like RT-PCR, the rapid antigen detection test too seeks to detect the virus rather than the [antibodies](#) produced by the body.
 - While the mechanism is different, the **most significant difference between the two is time**.
 - As the ICMR has pointed out, the **RT-PCR** test takes a minimum of **2-5 hours** including the time taken for sample transportation..
 - In a reliable **rapid antigen detection test**, the maximum duration for interpreting a positive or negative test is **30 minutes**.

RT-PCR Test

- **Kary Mullis**, the American biochemist invented the PCR technique. He was awarded the [Nobel Prize](#) for Chemistry in 1993.
- Under this, **copies of a segment of DNA (deoxyribonucleic acid)** are created using an enzyme called Polymerase.
 - The 'chain reaction' signifies how the DNA fragments are copied, exponentially — one is copied into two, the two are copied into four, and so on.
- A fluorescent **DNA binding dye called the “probe” is added to DNA**, which shows the presence of the virus on a [fluorometer](#).
- However, coronavirus is made of **RNA (ribonucleic acid)**.
- Therefore to detect coronavirus, **RNA is converted into DNA** using a technique called reverse transcription.
 - A 'reverse transcriptase' enzyme converts the RNA into DNA.
- Copies of the DNA are then made and amplified.

[Source: TH](#)

PDF Refernece URL: <https://www.drishtiiias.com/printpdf/radt-vs-rt-pcr>