

P Ovale Malaria

Why in News

Recently a **not very common type** of malaria, *Plasmodium ovale*, has been identified in a jawan in Kerala.

• The soldier is believed to have contracted it in **Sudan**, where Plasmodium ovale is endemic.

Key Points

About:

- Plasmodium ovale is one among the five kinds of malarial parasites Plasmodium falciparum, Plasmodium vivax (the commonest ones), Plasmodium Malariae, Plasmodium Ovale and Plasmodium Knowlesi.
- It is termed ovale as about 20% of the parasitised cells are oval in shape.
- The parasite can remain in the spleen or liver of the body for a long time, even years, after the mosquito bite, and the person could become symptomatic later.

Symptoms:

 Symptoms include fever for 48 hours, headache and nausea, and it rarely causes severe illness.

Similar to P vivax:

- P ovale is very similar to P vivax and the treatment modality is the same as it is for a person infected with P vivax.
- Distinguishing between P vivax and P ovale may be tricky and can be differentiated only through careful detection.

Prevalence:

- P ovale malaria is **endemic to tropical Western Africa.** It is relatively unusual outside of Africa and, where found, comprises less than 1% of the isolates.
- It has also been detected in the Philippines, Indonesia and Papua New Guinea, but is still relatively rare in these areas.


Transmission in India:

- According to the National Institute of Malaria Research (NIMR), the Kerala case could be an isolated one and there are no recorded cases of local transmission so far.
- Previously, too, isolated cases were reported in Gujarat, Kolkata, Odisha and Delhi.
 However, no local transmission has been recorded which means these cases have been acquired.
- In India, out of 1.57 lakh malaria cases in the high-burden states of Odisha,
 Chhattisgarh, Jharkhand, Meghalaya and Madhya Pradesh in 2019, 1.1 lakh cases (70%) were cases of falciparum malaria.
- According to the recent **World Malaria Report 2020**, cases in India dropped from about 20 million in 2000 to about 5.6 million in 2019.

Malaria

- Malaria is caused by Plasmodium (a protozoan).
- Malaria caused by **Plasmodium falciparum** is the **most serious** one and can even be **fatal.**
- Life Cycle of Plasmodium:
 - Plasmodium enters the human body as **sporozoites (infectious form)** through the bite of infected **female Anopheles mosquito.**
 - The parasites initially multiply within the liver cells and then attack the Red Blood Cells (RBCs) resulting in their rupture.
 - The rupture of RBCs is associated with release of a **toxic substance**, **haemozoin**, which is responsible for the chill and high fever recurring every three to four days.
 - When a female Anopheles mosquito bites an infected person, these parasites enter the mosquito's body and undergo further development.
 - The parasites multiply within them to form sporozoites that are stored in their salivary glands. When these mosquitoes bite a human, the sporozoites are introduced into his/ her body, thereby initiating the events mentioned above.





Note

- It is interesting to note that the malarial parasite requires two hosts human and mosquitoes
 to complete its life cycle.
- The female Anopheles mosquito is the vector (transmitting agent) too.
- World Malaria Day is observed on 25th April.
 - It can be noted that the World Health Organisation (WHO) officially endorses diseasespecific global awareness days for only four diseases viz. <u>HIV-AIDS</u>, <u>TB</u>, <u>Malaria</u>, <u>and</u> <u>Hepatitis</u>.

Source: IE

PDF Refernece URL: https://www.drishtiias.com/printpdf/p-ovale-malaria

